

Parathyroidectomy 1/2 (2 pages)

Brief description

Parathyroidectomy is surgery to remove parathyroid gland/s or parathyroid tumours'. Alternative name: - parathyroid gland removal the parathyroid glands are right behind your thyroid gland in the neck. These glands help the body control calcium.

Why is this operation necessary?

Parathyroidectomy is commonly recommended when the parathyroid glands are producing too much parathyroid hormone. This condition is called hyperparathyroidism. It is often caused by a small non-cancerous (benign) tumour called an adenoma.

Are there alternative treatments available?

Many factors are taken into consideration before deciding whether surgery is necessary and also what type. Some of these factors include your age, calcium levels and whether you have symptoms.

Is it safe to have this operation?


Before you agree to the operation, you should consider the risks that may be involved. Your surgery will be performed by a team of highly qualified and skilled professionals who will take all steps necessary to ensure a safe procedure and a successful result. However, there are risks involved with all surgery even if these risks may be very small.

What are the specific risks involved?

Hypoparathyroidism. This can lead to low calcium levels that are dangerous for your health. Injury to the thyroid gland or the need to remove part of the thyroid gland. Other complications may include: bleeding, a hoarse voice, difficulty swallowing, numbness of the skin on the neck, vocal cord paralysis. Occasionally the wound swells due to a build-up of blood in the neck within 24 hours after the operation. Rarely the wound needs to be re-examined in theatre. Chest infections may arise particularly in smokers. Wound infection is a rare problem and settles down with appropriate treatment. Aches and tingles may be felt in the wound for up to 6 months. Occasionally there may be numb patches in the skin around the wound, which gets better after 2 to 3 months.

What are the general risks involved?

There are risks for developing complications which are general and which may occur with any surgical procedure. These complications include the risk of infection, bleeding, pain, wound breakdown, deep vein thrombosis (blood clot in the legs that may travel to the lungs), or complications affecting the heart, lungs or kidneys.



What are the anaesthetic risks involved?

You can discuss the type of anaesthetic you will have with your anaesthetist and also the possible complications that may occur.

What should I do before the operation?

You should not eat or drink anything for at least 8 hours before your operation. However, you should take all your regular medication as usual on the day. You will need to stop certain medication such as disprin, warfarin, clopidogrel, plavix or other blood thinning medicines (including homeopathic medication) well before the operation. Please discuss with your surgeon. This is often a busy period. You may require further investigations to assess your fitness for surgery.

How long does the operation take?

Usually about 1 to 2 hours

What happens when I wake up?

After the operation is completed, you will be transferred back to the ward. Although you will be conscious a minute or two after the operation ends, you are unlikely to remember anything until you are back in your bed on the ward. You will have a drip in your arm and a drain in your neck.

Will I have pain?

Some pain may be present, but this should be controlled to a level of mild discomfort with the painkillers that are prescribed. Ask the nursing staff for medication if you have pain.

What happens before the operation?

Please report to the hospital reception on time for your admission. Please bring along all the documents that may be required such as your medical aid card, id and contact details. If you are not a member of a medical aid you will be required to pay a deposit or to sign an indemnity form. As far as possible we will try to advise you about hospital costs before your admission. It may be best to complete some of the documentation beforehand at the hospital pre-admission clinic to save time on the day of your admission. When you arrive in the ward, you will be welcomed by the nurses or the receptionist and will have your details checked. Some basic tests will be done such as pulse, temperature, blood pressure and urine examination. You will be asked to hand in any medicines or drugs you may be taking, so that your drug treatment in hospital will be correct. Please tell the nurses of any allergies to drugs or dressings. The surgeon will have explained the operation and you will be asked to sign your consent for the operation. If you are not clear about any part of the operation, ask for more details from the surgeon or from the nurses. In an adult the operation area may need to be shaved to remove excess hair. You may be issued with compression stockings that will help prevent blood clots in your legs. If you are having a general anaesthetic, the anaesthetist who will be giving your anaesthetic will interview and examine you and he may put up a drip or prescribe some medication to help you relax. You will be taken on a trolley to the operating suite by the staff. You will be wearing a cotton gown. Wedding rings will be fastened with tape and removable dentures will be left on the ward. It is best to leave all jewellery at home. There will be several checks on your details on the way to the operating theatre where your anaesthetic will begin.

How soon after the operation can I eat?

You will be allowed water, tea or juice in small amounts immediately after the operation, but may only start eating food again when fully awake.

How soon after the operation can I get out of bed?

You should be able to walk a short distance very soon after waking up, but ask the nursing staff for assistance if you feel dizzy. You should be able to walk without too much discomfort by the next day and will be encouraged to mobilize as much as possible.

How long will I stay in the hospital?

Usually, 1 to 2 days

What happens when I am discharged from the ward?

Your surgeon will determine when you are ready to go home. You will be given some medication for pain. You will be given instructions on the dressings and how to care for the wound. You will also get an appointment for your follow-up in the surgeon's rooms. You should ask for a sick certificate if you need this for your employer.

What should I be aware of when I get home?

Some symptoms may not be evident for 2 to three days after surgery. If you experience any of the following, call your surgeon. Increasing pain, fever, swelling, wound discharge, shortness of breath, numbness and tingling around the lips and hands.

How soon can I start exercise?

You can perform routine activities as soon as you get home. Mild exercise like walking or climbing stairs would be possible within the first day or 2 and full exercise after 4 weeks.

How soon can I drive a car?

You can drive as soon as you can make an emergency stop and look around in all directions without hurting your neck. Usually within a week.

How long will I be off work?

You should be able to return to a light job within 2 weeks and any heavy job within 4 weeks

What about payment?

The procedure and its associated costs will have been discussed with you, and a quote provided. Where procedures need to be unexpectedly altered during the course of the procedure, the fee may change. Similarly, emergency procedures may incur an additional cost. Please settle your account promptly.